


Highlights of the 2010 American Heart Association Guidelines for CPR & ECC



Summary of Key BLS Components for Adults, Children, and Infants*

	Recommendations		
Component	Adults	Children	Infants
Recognition	Unresponsive (for all ages)		
	No breathing or no normal breathing (ie, only gasping)	No breathing or only gasping	
	No pulse palpated within 10 seconds for all ages (HCP only)		
CPR sequence	C-A-B		
Compression rate	At least 100/min		
Compression depth	At least 2 inches (5 cm)	At least ⅓ AP diameter About 2 inches (5 cm)	At least ⅓ AP diameter About 1½ inches (4 cm)
Chest wall recoil	Allow complete recoil between compressions HCPs rotate compressors every 2 minutes		
Compression interruptions	Minimize interruptions in chest compressions Attempt to limit interruptions to <10 seconds		
Airway	Head tilt–chin lift (HCP suspected trauma: jaw thrust)		
Compression-to-ventilation ratio (until advanced airway placed)	30:2 1 or 2 rescuers	30:2 Single rescuer 15:2 2 HCP rescuers	
Ventilations: when rescuer untrained or trained and not proficient	Compressions only		
Ventilations with advanced airway (HCP)	1 breath every 6-8 seconds (8-10 breaths/min) Asynchronous with chest compressions About 1 second per breath Visible chest rise		
Defibrillation	Attach and use AED as soon as available. Minimize interruptions in chest compressions before and after shock; resume CPR beginning with compressions immediately after each shock.		

Abbreviations: AED, automated external defibrillator; AP, anterior-posterior; CPR, cardiopulmonary resuscitation; HCP, healthcare provider.

*Excluding the newly born, in whom the etiology of an arrest is nearly always asphyxial.

Reprinted from *Highlights of the 2010 AHA Guidelines for CPR & ECC*; <http://static.heart.org/eccguidelines/guidelines-highlights.html>; copyright 2010.